

Revised version for publication in Israel J. Math.

EXTERIOR ALGEBRAS AND TWO CONJECTURES ON FINITE ABELIAN GROUPS

TAO FENG¹, ZHI-WEI SUN² AND QING XIANG³

ABSTRACT. Let G be a finite abelian group with $|G| > 1$. Let a_1, \dots, a_k be k distinct elements of G and let b_1, \dots, b_k be (not necessarily distinct) elements of G , where k is a positive integer smaller than the smallest prime divisor of $|G|$. We show that there is a permutation π on $\{1, \dots, k\}$ such that $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct, provided that any other prime divisor of $|G|$ (if there is any) is greater than $k!$. This in particular confirms the Dasgupta-Károlyi-Serra-Szegedy conjecture for abelian p -groups. We also pose a new conjecture involving determinants and characters, and show that its validity implies Snevily's conjecture for abelian groups of odd order. Our methods involve exterior algebras and characters.

1. INTRODUCTION

Let $G = \{a_1, \dots, a_n\}$ be an abelian group (written multiplicatively) of order n , and let $b_1, \dots, b_n \in G$. In 1952, M. Hall, Jr. [7] showed that $a_1 b_{\pi(1)}, \dots, a_n b_{\pi(n)}$ are (pairwise) distinct for some permutation $\pi \in S_n$ (the symmetric group on $\{1, \dots, n\}$) if and only if $b_1 \cdots b_n$ is the identity element of G .

In 1999, H. S. Snevily [9] considered subsets with cardinality k of an abelian group G (or, simply k -subsets of G), and proposed the following challenging conjecture.

Conjecture 1.1. (Snevily's Conjecture) *Let G be a multiplicatively written abelian group of odd order, and let $A = \{a_1, \dots, a_k\}$ and $B = \{b_1, \dots, b_k\}$ be two k -subsets of G . Then there is a permutation $\pi \in S_k$ such that $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct.*

The above conjecture can be reformulated in terms of Latin transversals of the Latin square formed by the Cayley multiplication table of G . N. Alon [2] proved Snevily's conjecture when $|G|$ is an odd prime, by using the Combinatorial Nullstellensatz [1]. In fact, Alon [2] obtained the following stronger result.

Theorem 1.2. (Alon [2]) *Let G be a cyclic group of prime order p . Let $k < p$ be a positive integer. Let $A = \{a_1, \dots, a_k\}$ be a k -subset of G and b_1, \dots, b_k be (not necessarily distinct) elements of G . Then there is a permutation $\pi \in S_k$ such that $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct.*

Key words and phrases. Abelian group, additive combinatorics, exterior algebra, character, Latin square.

2000 *Mathematics Subject Classification.* Primary 20D60; Secondary 05A05, 05E99, 11B75, 11P99, 15A75, 20K01.

¹Research supported in part by National Natural Science Foundation of China (grant 10331030).

²Corresponding author. Supported by National Natural Science Foundation of China (grant 10871087).

³Research supported in part by NSF Grant DMS 0701049.

In 2001, S. Dasgupta, G. Károlyi, O. Serra and B. Szegedy [4] proved Conjecture 1.1 for any cyclic group G of odd order. Moreover, these authors extended Alon's result (Theorem 1.2) to cyclic groups of prime power order as well as elementary abelian groups.

Theorem 1.3. (Dasgupta-Károlyi-Serra-Szegedy [4]) *Let p be a prime and let α be a positive integer. Let G be the cyclic group C_{p^α} or the elementary abelian p -group C_p^α . Assume that $A = \{a_1, a_2, \dots, a_k\}$ is a k -subset of G and b_1, b_2, \dots, b_k are (not necessarily distinct) elements of G , where k is smaller than p . Then, for some $\pi \in S_k$ the products $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct.*

Motivated by Theorems 1.2 and 1.3, Dasgupta, et al. [4, p. 23] proposed the following conjecture.

Conjecture 1.4. (The DKSS Conjecture) *Let G be a finite abelian group with $|G| > 1$, and let $p(G)$ be the smallest prime divisor of $|G|$. Let $k < p(G)$ be a positive integer. Assume that $A = \{a_1, a_2, \dots, a_k\}$ is a k -subset of G and b_1, b_2, \dots, b_k are (not necessarily distinct) elements of G . Then there is a permutation $\pi \in S_k$ such that $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct.*

Let G be a finite abelian group with $|G| > 1$. When $A = \{a_1, \dots, a_k\}$ is a subgroup of G with cardinality k and b_1, \dots, b_k are (not necessarily distinct) elements of A , by Hall's theorem, $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct for some $\pi \in S_k$ if and only if $b_1 \cdots b_k = e$. As G always has an element of order $p = p(G)$, which generates a cyclic subgroup of order p , we see that the conclusion of Conjecture 1.4 does not hold in general when $k = p(G)$. We also note that Conjecture 1.4 implies Snevily's conjecture in the case where $k < p(G)$. Using a group ring approach, W. D. Gao and D. J. Wang [5] proved Conjecture 1.4 for abelian p -groups under the stronger assumption $k < \sqrt{2p}$. (See also [GGH].)

In this paper we confirm the DKSS conjecture under the extra assumption that the second smallest prime divisor of $|G|$ (if it exists) is greater than $k!$. It will be convenient to introduce the following terminology.

Definition 1.5. Let k and $n > 1$ be positive integers. We say that n is k -large if the smallest prime divisor of n is greater than k and any other prime divisor of n is greater than $k!$.

Here is our main result on the DKSS conjecture.

Theorem 1.6. *Let G be a finite abelian group. Let $A = \{a_1, \dots, a_k\}$ be a k -subset of G , and b_1, \dots, b_k be (not necessarily distinct) elements of G . Suppose that either A or $B = \{b_1, \dots, b_k\}$ is contained in a subgroup H of G and $|H|$ is k -large. Then there exists a permutation $\pi \in S_k$ such that $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct.*

Note that if k is a positive integer and p is a prime such that $p > k$, then p^α is obviously k -large for every $\alpha = 1, 2, 3, \dots$. Therefore we have the following immediate consequence of Theorem 1.6.

Corollary 1.7. *Let p be a prime. Assume that G is an abelian p -group, and k is a positive integer such that $k < p$. Let $A = \{a_1, \dots, a_k\}$ be a k -subset of G , and b_1, b_2, \dots, b_k be (not necessarily distinct) elements of G . Then there is a permutation $\pi \in S_k$ such that $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct.*

Obviously Theorem 1.6 implies that the DKSS conjecture is true for $k = 3$. By a case-by-case analysis we also can show that the DKSS conjecture holds when $k = 4$. Theorem 1.6 also implies that the DKSS conjecture holds in the case where $A = \{a, a^2, \dots, a^k\}$ with $a \in G \setminus \{e\}$.

Corollary 1.8. *Let G be a finite abelian group with $|G| > 1$. Let $a \neq e$ be an element of G and let b_1, \dots, b_k be (not necessarily distinct) elements of G . Provided that $k < p(G)$, there is a permutation $\pi \in S_k$ such that the products $a^i b_{\pi(i)}$ ($i = 1, \dots, k$) are distinct.*

It is interesting to compare Corollary 1.8 with the following conjecture of Snevily [9]: If $G = \langle a \rangle$ is a cyclic group of order n , k is a positive integer less than n , and b_1, \dots, b_k are (not necessarily distinct) elements of G , then there is a permutation $\pi \in S_k$ such that $a^i b_{\pi(i)}$, $i = 1, \dots, k$, are distinct.

Our proof of Theorem 1.6 uses an exterior algebra approach. For applications of symmetric product and alternating product methods in combinatorics, we refer the reader to [3, Chap. 6]. The exterior algebra approach can also be used to give an alternative proof of the following theorem of Z. W. Sun [14].

Theorem 1.9. (Sun [14]) *Let G be a finite cyclic group, and let $A_1 = \{a_{11}, \dots, a_{1k}\}$, $A_2 = \{a_{21}, \dots, a_{2k}\}, \dots, A_m = \{a_{m1}, \dots, a_{mk}\}$ be k -subsets of G , where $m \in \{3, 5, 7, \dots\}$. Then there exist permutations $\pi_i \in S_k$, $2 \leq i \leq m$, such that $a_{1j} a_{2\pi_2(j)} \cdots a_{m\pi_m(j)}$, $j = 1, 2, \dots, k$, are all distinct.*

We remark that for $m = 2, 4, 6, \dots$, Sun could prove a similar result under the assumption that all elements of A_m have odd order (cf. [13] and [14]). This result in particular implies that Snevily's conjecture is true for cyclic groups of odd order, a result first proved by Dasgupta, et al. in [4].

Recall that, for a matrix $M = (a_{ij})_{1 \leq i, j \leq k}$ over a commutative ring with identity, the determinant of M and the permanent of M are defined by

$$\det(M) = \sum_{\sigma \in S_k} \text{sgn}(\sigma) a_{1\sigma(1)} \cdots a_{k\sigma(k)} \text{ and } \text{per}(M) = \sum_{\sigma \in S_k} a_{1\sigma(1)} \cdots a_{k\sigma(k)}$$

respectively, where $\text{sgn}(\sigma)$, the sign of $\sigma \in S_k$, equals 1 or -1 according as the permutation σ is even or odd.

To attack the Snevily conjecture via our approach, we propose the following conjecture.

Conjecture 1.10. *Let G be a finite abelian group, and let $A = \{a_1, \dots, a_k\}$ and $B = \{b_1, \dots, b_k\}$ be two k -subsets of G . Let K be any field containing an element of multiplicative order $|G|$, and let \hat{G} be the character group of all group homomorphisms from G to $K^* = K \setminus \{0\}$. Then there are $\chi_1, \dots, \chi_k \in \hat{G}$ such that $\det(\chi_i(a_j))_{1 \leq i, j \leq k}$ and $\det(\chi_i(b_j))_{1 \leq i, j \leq k}$ are both nonzero.*

When G is cyclic, we may take $\chi_i = \chi^i$ for $i = 1, 2, \dots, k$, where χ is a generator of \hat{G} . Then the two determinants $\det(\chi_i(a_j))_{1 \leq i, j \leq k}$ and $\det(\chi_i(b_j))_{1 \leq i, j \leq k}$ in the above conjecture are both nonzero since they are Vandermonde determinants. Therefore we see that Conjecture 1.10 is true for cyclic groups. We further mention that when G is a cyclic group of prime order and K is the complex field \mathbb{C} , for any distinct $\chi_1, \dots, \chi_k \in \hat{G}$ and distinct $a_1, \dots, a_k \in G$ we have $\det(\chi_i(a_j))_{1 \leq i, j \leq k} \neq 0$ by the Chebotarëv theorem (cf.

[10] and [15]); this is stronger than what Conjecture 1.10 asserts. The general case of Conjecture 1.10 seems to be quite sophisticated.

We will show that Conjecture 1.10 holds when $A = B$ (see Lemma 3.1). Moreover, we have the following result.

Theorem 1.11. (i) *Conjecture 1.10 implies Conjecture 1.1.*

(ii) *Let G, A, B, \hat{G} be as in Conjecture 1.10. Assume that there is a $\pi \in S_k$ such that $C = \{a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}\}$ is a k -set with $\{a_1 b_{\tau(1)}, \dots, a_k b_{\tau(k)}\} \neq C$ for all $\tau \in S_k \setminus \{\pi\}$. Then there are $\chi_1, \dots, \chi_k \in \hat{G}$ such that $\det(\chi_i(a_j))_{1 \leq i, j \leq k}$ and $\det(\chi_i(b_j))_{1 \leq i, j \leq k}$ are both nonzero. Also, there are $\chi_1, \dots, \chi_k \in \hat{G}$ such that $\det(\chi_i(a_j))_{1 \leq i, j \leq k}$ and $\text{per}(\chi_i(b_j))_{1 \leq i, j \leq k}$ are both nonzero, and there are $\psi_1, \dots, \psi_k \in \hat{G}$ such that the permanents $\text{per}(\psi_i(a_j))_{1 \leq i, j \leq k}$ and $\text{per}(\psi_i(b_j))_{1 \leq i, j \leq k}$ are both nonzero.*

In the next section we will give some background on exterior algebras and lay the basis for our new approach. (For readers who are not familiar with exterior algebras, Northcott's book [8] is a good reference.) We are going to prove Theorem 1.6 and Corollary 1.8 in Section 3, and Theorems 1.9 and 1.11 in Section 4.

2. AN AUXILIARY PROPOSITION MOTIVATED BY EXTERIOR ALGEBRAS

Let us recall some definitions and basic facts related to exterior algebras.

Let R be a commutative ring with identity, and let M be a left R -module. The n th exterior power of M , denoted by $\bigwedge^n M$, comes equipped with an alternating multilinear map

$$M \times M \times \cdots \times M \rightarrow \bigwedge^n M, (m_1, m_2, \dots, m_n) \mapsto m_1 \wedge m_2 \wedge \cdots \wedge m_n,$$

that is universal: for an R -module N and an alternating multilinear map $\beta : M \times M \times \cdots \times M \rightarrow N$, there is a unique linear map from $\bigwedge^n M$ to N which takes $m_1 \wedge m_2 \wedge \cdots \wedge m_n$ to $\beta(m_1, m_2, \dots, m_n)$. Recall that a multilinear map β is *alternating* if $\beta(m_1, m_2, \dots, m_n) = 0$ whenever two of the m_i are equal. The exterior power $\bigwedge^n M$ can be constructed as the quotient module of $M^{\otimes n}$ (the n -th tensor power) by the submodule generated by all those $m_1 \otimes m_2 \otimes \cdots \otimes m_n$ with two of the m_i equal. We naturally identify $\bigwedge^0 M = R$ and $\bigwedge^1 M = M$. The *exterior algebra* of M , denoted by $E(M)$, is the algebra $\bigoplus_{n \geq 0} \bigwedge^n(M)$, with respect to the wedge product ' \wedge '. This is a graded algebra. By 'graded' we mean that multiplying an element of $\bigwedge^m M$ with an element of $\bigwedge^n M$, gives an element of $\bigwedge^{m+n} M$. A *skew derivation* on $E(M)$ is an R -homomorphism $\Delta : E(M) \rightarrow E(M)$ such that

$$\Delta(xy) = (\Delta x)y + (-1)^n x(\Delta y),$$

for all $x \in \bigwedge^n M$ and $y \in E(M)$.

Next let U be an R -module. Assume that we have a bilinear mapping $\gamma : U \times M \rightarrow R$. Then, for any $u \in U$, $\gamma(u, \cdot)$ is an R -module homomorphism from M to R . By [8, Theorem 10, p. 96], there exists a unique skew derivation $\Delta_u : E(M) \rightarrow E(M)$ that extends $\gamma(u, \cdot)$. Furthermore, when $n > 0$, Δ_u maps $\bigwedge^n M$ into $\bigwedge^{n-1} M$, and it can be defined by

$$\Delta_u(m_1 \wedge \cdots \wedge m_n) = \sum_{i=1}^n (-1)^{i+1} \gamma(u, m_i) (m_1 \wedge \cdots \wedge \widehat{m}_i \wedge \cdots \wedge m_n),$$

where m_1, \dots, m_n are arbitrary elements of M , and $m_1 \wedge \dots \wedge \widehat{m_i} \wedge \dots \wedge m_n$ denotes the result of striking out m_i from $m_1 \wedge \dots \wedge m_i \wedge \dots \wedge m_n$. For $u_1, u_2 \in U$, we can consider the composition $\Delta_{u_1} \circ \Delta_{u_2}$ of the R -module homomorphisms Δ_{u_1} and Δ_{u_2} in the usual sense. We are now ready to state the following result from [8].

Lemma 2.1. ([8, Corollary, p. 100]) *Using the above notation, for $u_1, \dots, u_k \in U$ and $m_1, \dots, m_k \in M$, we have*

$$(\Delta_{u_1} \circ \dots \circ \Delta_{u_k})(m_1 \wedge \dots \wedge m_k) = (-1)^{k(k-1)/2} \det(\gamma(u_i, m_j))_{1 \leq i, j \leq k}.$$

Lemma 2.1 leads us to the following useful proposition.

Proposition 2.2. *Let G be a finite abelian group. Let \hat{G} denote the group of characters from G to $K^* = K \setminus \{0\}$, where K is a field containing an element of multiplicative order $|G|$. Let $a_1, \dots, a_k, b_1, \dots, b_k \in G$ and $\chi_1, \dots, \chi_k \in \hat{G}$. Suppose that both $\det(M_a)$ and $\text{per}(M_b)$ are nonzero, where $M_a = (\chi_i(a_j))_{1 \leq i, j \leq k}$ and $M_b = (\chi_i(b_j))_{1 \leq i, j \leq k}$. Then there is a permutation $\pi \in S_k$ such that the products $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct.*

Proof. For the purpose of applying Lemma 2.1, we set $R := K$, $M := KG$ (the group algebra of G over K), and $U := K\hat{G}$ (the group algebra of \hat{G} over K). The mapping $\gamma : U \times M \rightarrow K$ is defined as follows: First define $\gamma : \hat{G} \times G \rightarrow K$ by setting $\gamma(\chi, g) := \chi(g)$ for $\chi \in \hat{G}$ and $g \in G$; next we extend γ bilinearly to a map from $U \times M$ to K . The resulting map is still denoted by γ and it is bilinear.

For any $\pi \in S_k$ we set

$$Q_\pi := a_1 b_{\pi(1)} \wedge \dots \wedge a_k b_{\pi(k)} \in \bigwedge^k M.$$

Let $M_{a,b}^\pi$ be the $k \times k$ matrix with (i, j) -entry equal to $\chi_i(a_j b_{\pi(j)})$. By Lemma 2.1, we have

$$\begin{aligned} (\Delta_{\chi_1} \circ \dots \circ \Delta_{\chi_k})(Q_\pi) &= (-1)^{k(k-1)/2} \det(M_{a,b}^\pi) \\ &= (-1)^{k(k-1)/2} \sum_{\sigma \in S_k} \text{sgn}(\sigma) \chi_1(a_{\sigma(1)} b_{\pi\sigma(1)}) \cdots \chi_k(a_{\sigma(k)} b_{\pi\sigma(k)}). \end{aligned} \quad (2.1)$$

Summing (2.1) over $\pi \in S_k$, we obtain

$$\begin{aligned} &(\Delta_{\chi_1} \circ \dots \circ \Delta_{\chi_k}) \left(\sum_{\pi \in S_k} Q_\pi \right) \\ &= (-1)^{k(k-1)/2} \sum_{\sigma \in S_k} \sum_{\pi \in S_k} \chi_1(a_{\sigma(1)} b_{\pi\sigma(1)}) \cdots \chi_k(a_{\sigma(k)} b_{\pi\sigma(k)}) \text{sgn}(\sigma) \\ &= (-1)^{k(k-1)/2} \sum_{\sigma \in S_k} \text{sgn}(\sigma) \chi_1(a_{\sigma(1)}) \cdots \chi_k(a_{\sigma(k)}) \sum_{\pi \in S_k} \chi_1(b_{\pi(1)}) \cdots \chi_k(b_{\pi(k)}) \\ &= (-1)^{k(k-1)/2} \det(M_a) \text{per}(M_b). \end{aligned} \quad (2.2)$$

As $\det(M_a) \text{per}(M_b) \neq 0$, we have $\sum_{\pi \in S_k} Q_\pi \neq 0$. So there exists a $\pi \in S_k$ such that $a_1 b_{\pi(1)} \wedge \dots \wedge a_k b_{\pi(k)}$ is nonzero, implying that $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct. The proof is now complete.

3. PROOFS OF THEOREM 1.6 AND COROLLARY 1.8

Lemma 3.1. *Let G be a finite abelian group and let K be a field containing an element of multiplicative order $|G|$. Let \hat{G} be the group of characters from G to K^* , and let $a_1, \dots, a_k \in G$. Then a_1, \dots, a_k are distinct if and only if there are (distinct) $\chi_1, \dots, \chi_k \in \hat{G}$ such that $\det(\chi_i(a_j))_{1 \leq i, j \leq k} \neq 0$. Also, there exist $\chi_1, \dots, \chi_k \in \hat{G}$ with $\text{per}(\chi_i(a_j))_{1 \leq i, j \leq k} \neq 0$ provided that a_1, \dots, a_k are distinct.*

Proof. If $a_s = a_t$ for some $1 \leq s < t \leq k$, then for any $\chi_1, \dots, \chi_k \in \hat{G}$ the determinant $\det(\chi_i(a_j))_{1 \leq i, j \leq k}$ vanishes since the s th column and t th column of the matrix $(\chi_i(a_j))_{1 \leq i, j \leq k}$ are identical.

Now suppose that a_1, \dots, a_k are distinct. If the characteristic of K is a prime p dividing $|G|$, then

$$(x^{|G|/p} - 1)^p = x^{|G|} - 1 \quad \text{for all } x \in K,$$

which contradicts the assumption that K contains an element of multiplicative order $|G|$. So we have $|G|1 \neq 0$, where 1 is the identity of the field K . It is well known that

$$\sum_{\chi \in \hat{G}} \chi(a) = \begin{cases} 0, & \text{if } a \in G \setminus \{e\}, \\ |G|1, & \text{if } a = e. \end{cases}$$

Observe that

$$\begin{aligned} & \sum_{\chi_1, \dots, \chi_k \in \hat{G}} \chi_1(a_1^{-1}) \cdots \chi_k(a_k^{-1}) \det(\chi_i(a_j))_{1 \leq i, j \leq k} \\ &= \sum_{\chi_1, \dots, \chi_k \in \hat{G}} \chi_1(a_1^{-1}) \cdots \chi_k(a_k^{-1}) \sum_{\pi \in S_k} \text{sgn}(\pi) \prod_{i=1}^k \chi_i(a_{\pi(i)}) \\ &= \sum_{\chi_1, \dots, \chi_k \in \hat{G}} \sum_{\pi \in S_k} \text{sgn}(\pi) \prod_{i=1}^k \chi_i(a_{\pi(i)} a_i^{-1}) \\ &= \sum_{\pi \in S_k} \text{sgn}(\pi) \prod_{i=1}^k \sum_{\chi_i \in \hat{G}} \chi_i(a_{\pi(i)} a_i^{-1}) \\ &= \text{sgn}(I) \prod_{i=1}^k (|G|1) = (|G|1)^k \neq 0, \end{aligned}$$

where I is the identity permutation in S_k . So there are $\chi_1, \dots, \chi_k \in \hat{G}$ such that $\det(\chi_i(a_j))_{1 \leq i, j \leq k} \neq 0$ and hence χ_1, \dots, χ_k are distinct. Similarly, by removing $\text{sgn}(\pi)$ in the above calculations, we see that there exist $\chi_1, \dots, \chi_k \in \hat{G}$ such that $\text{per}(\chi_i(a_j))_{1 \leq i, j \leq k} \neq 0$. This concludes the proof.

Remark 3.2. If we apply Lemma 3.1 with $k = |G|$ then we obtain the following classical result: The matrix $T = (\chi(g))_{\chi \in \hat{G}, g \in G}$ is nonsingular; in other words, all the characters in \hat{G} are linearly independent over the field K . It is well known that all the characters in

\hat{G} actually form a basis of the vector space

$$K^G = \{f : f \text{ is a function from } G \text{ to } K\}$$

over the field K .

Lemma 3.3. (Sun [12, Lemma 3.1]) *Let $\lambda_1, \dots, \lambda_k$ be complex n th roots of unity. Suppose that $c_1\lambda_1 + \dots + c_k\lambda_k = 0$, where c_1, \dots, c_k are nonnegative integers. Then $c_1 + \dots + c_k$ can be written in the form $\sum_{p|n} px_p$, where the sum is over all prime divisors of n and the x_p are nonnegative integers.*

This lemma appeared in [12] explicitly, and it follows from Lemma 9 in Sun [11].

We are now in a position to prove Theorem 1.6.

Proof of Theorem 1.6. Let \hat{G} be the group of all complex-valued characters of G , and let $p_1 < \dots < p_r$ be all the distinct prime divisors of $|H|$. If $k! = p_1x_1 + \dots + p_rx_r$ for some nonnegative integers x_1, \dots, x_r , then $k! = p_ix_i$ for some $1 \leq i \leq r$ (since $p_{i_1} + p_{i_2} > k!$ for all $1 \leq i_1 < i_2 \leq r$), thus p_i divides $k!$, contradicting the fact $k < p(H) \leq p_i$. In view of this and Lemma 3.3, we see that if $\zeta_\pi \in \mathbb{C}$ and $\zeta_\pi^{|H|} = 1$ for all $\pi \in S_k$ then $\sum_{\pi \in S_k} \zeta_\pi \neq 0$.

Case 1. $B = \{b_1, \dots, b_k\} \subseteq H$.

In this case $b_1^{|H|} = \dots = b_k^{|H|} = e$. As a_1, \dots, a_k are distinct, by Lemma 3.1 there are $\chi_1, \dots, \chi_k \in \hat{G}$ such that $\det(\chi_i(a_j))_{1 \leq i, j \leq k} \neq 0$. Observe that

$$\text{per}(\chi_i(b_j))_{1 \leq i, j \leq k} = \sum_{\pi \in S_k} \prod_{i=1}^k \chi_i(b_{\pi(i)})$$

and

$$\left(\prod_{i=1}^k \chi_i(b_{\pi(i)}) \right)^{|H|} = \prod_{i=1}^k \chi_i(b_{\pi(i)}^{|H|}) = \prod_{i=1}^k \chi_i(e) = 1.$$

By the previous discussion, $\text{per}(\chi_i(b_j))_{1 \leq i, j \leq k} \neq 0$. It follows from Proposition 2.2 that $a_1b_{\pi(1)}, \dots, a_kb_{\pi(k)}$ are distinct for some $\pi \in S_k$.

Case 2. $A \subseteq H$.

Suppose that $p_1, \dots, p_r, \dots, p_s$ ($s \geq r$) are all the distinct prime divisors of $|G|$. It is well known that $G = P_1 \cdots P_r \cdots P_s$, where each $P_i = \text{Syl}_{p_i}(G)$ is the unique Sylow p_i -subgroup of G . For each $i = 1, \dots, r$, we have $\text{Syl}_{p_i}(H) \subseteq P_i$ by Sylow's second theorem. So

$$H = \text{Syl}_{p_1}(H) \cdots \text{Syl}_{p_r}(H) \subseteq H_1 := P_1 \cdots P_r.$$

As a_1, \dots, a_k are distinct elements of H_1 , by Lemma 3.1 there are $\chi_1, \dots, \chi_k \in \widehat{H_1}$ such that $\det(\chi_i(a_j))_{1 \leq i, j \leq k} \neq 0$. For each $i = 1, \dots, k$ we can extend χ_i to a character of G by setting $\chi_i(h_1h_2) = \chi_i(h_1)$ for all $h_1 \in H_1$ and $h_2 \in H_2$, where $H_2 = P_{r+1} \cdots P_s$ if $r < s$, and $H_2 = \{e\}$ if $r = s$. It follows that $\chi_i(g)^{|H_1|} = 1$ for all $g \in G = H_1H_2$. Thus

$$\left(\prod_{i=1}^k \chi_i(b_{\pi(i)}) \right)^{|H_1|} = 1 \quad \text{for all } \pi \in S_k.$$

Note that $|H_1|$ is k -large since $|H|$ is. As in Case 1, we get $\text{per}(\chi_i(b_j))_{1 \leq i, j \leq k} \neq 0$ and hence the desired result follows.

Combining the above, we have completed the proof of Theorem 1.6.

Proof of Corollary 1.8. Let p be any prime divisor of the order of a , and let $P = \text{Syl}_p(G)$, the Sylow p -subgroup of G . As the order of a does not divide $|G|/|P|$, we have $\tilde{a} = a^{|G|/|P|} \in P \setminus \{e\}$.

The condition $k < p(G)$ ensures that $1, \dots, k$ are pairwise incongruent modulo p . The following argument actually yields a refinement of the stated result.

Let i_1, i_2, \dots, i_k be any integers pairwise incongruent modulo p . Then $\tilde{a}^{i_1}, \tilde{a}^{i_2}, \dots, \tilde{a}^{i_k}$ are distinct elements of P . Set $\tilde{b}_j = b_j^{|G|/|P|}$ for $j = 1, \dots, k$. By Theorem 1.6, there is a permutation $\pi \in S_k$ such that all the elements

$$\tilde{a}^{i_j} \widetilde{b_{\pi(j)}} = (a^{i_j} b_{\pi(j)})^{|G|/|P|} \quad (j = 1, \dots, k)$$

are distinct. It follows that $a^{i_1} b_{\pi(1)}, a^{i_2} b_{\pi(2)}, \dots, a^{i_k} b_{\pi(k)}$ are distinct. We are done.

4. PROOFS OF THEOREMS 1.9 AND 1.11

First, we prove Theorem 1.9 via the exterior algebra approach.

Proof of Theorem 1.9. We choose $K = \mathbb{C}$ (the field of complex numbers), and work with the group algebra $M := KG$. For $\pi_2, \dots, \pi_m \in S_k$ we define

$$Q_{\pi_2, \dots, \pi_m} = \prod_{i=2}^m \text{sgn}(\pi_i) (a_{11} a_{2\pi_2(1)} \cdots a_{m\pi_m(1)} \wedge \cdots \wedge a_{1k} a_{2\pi_2(k)} \cdots a_{m\pi_m(k)}) \in \bigwedge^k M.$$

Let χ_1, \dots, χ_k be distinct (complex) characters of G , and set $M_i = (\chi_\ell(a_{ij}))_{1 \leq \ell, j \leq k}$ for $i = 1, 2, \dots, m$. Then, by similar computations to those in (2.1) and (2.2) (paying attention to the signs involved and noting that m is odd), we get

$$\Delta_{\chi_1} \circ \cdots \circ \Delta_{\chi_k} \left(\sum_{\pi_2, \dots, \pi_m \in S_k} Q_{\pi_2, \dots, \pi_m} \right) = (-1)^{k(k-1)/2} \prod_{i=1}^m \det(M_i).$$

Since G is cyclic, we have $\hat{G} = \{1, \chi, \chi^2, \dots\}$, where χ is a generator of \hat{G} . If $\chi(a_{is}) = \chi(a_{it})$, then $\sum_{\psi \in \hat{G}} \psi(a_{is} a_{it}^{-1}) = |G|1 \neq 0$, hence $a_{is} a_{it}^{-1} = e$ and thus $s = t$ (since a_{i1}, \dots, a_{ik} are distinct). Therefore, if we choose $\chi_\ell = \chi^{\ell-1}$ for $\ell = 1, 2, \dots, k$, then $\det(M_i) \neq 0$ for all $1 \leq i \leq m$ since each M_i is a Vandermonde matrix whose second row is $(\chi(a_{ij}))_{1 \leq j \leq k}$. With the above choice of χ_ℓ , $1 \leq \ell \leq k$, we have

$$\Delta_{\chi_1} \circ \cdots \circ \Delta_{\chi_k} \left(\sum_{\pi_2, \dots, \pi_m \in S_k} Q_{\pi_2, \dots, \pi_m} \right) \neq 0.$$

Hence there exist $\pi_2, \dots, \pi_m \in S_k$ such that $Q_{\pi_2, \dots, \pi_m} \neq 0$, implying that $a_{1j} a_{2\pi_2(j)} \cdots a_{m\pi_m(j)}$, $1 \leq j \leq k$, are all distinct.

Next, we give a proof of Theorem 1.11.

Proof of Theorem 1.11. (i) We want to prove Conjecture 1.1 under the assumption that Conjecture 1.10 holds.

As $|G|$ is odd, we have $2^{\varphi(|G|)} \equiv 1 \pmod{|G|}$ by Euler's theorem, where φ is the Euler totient function. Let K be the finite field $\mathbb{F}_{2^{\varphi(|G|)}}$. Then the cyclic group $K^* = K \setminus \{0\}$ has an element of multiplicative order $|G|$. Let \hat{G} be the group of characters from G to K^* . By Conjecture 1.10 there are $\chi_1, \dots, \chi_k \in \hat{G}$ such that $\det(M_a)$ and $\det(M_b)$ are both nonzero, where $M_a = (\chi_i(a_j))_{1 \leq i, j \leq k}$ and $M_b = (\chi_i(b_j))_{1 \leq i, j \leq k}$. As K is of characteristic 2, we have $\text{per}(M_b) = \det(M_b) \neq 0$. Applying Proposition 2.2 we obtain that $a_1 b_{\pi(1)}, \dots, a_k b_{\pi(k)}$ are distinct for some $\pi \in S_k$.

(ii) Set

$$\Sigma := \sum_{\chi_1, \dots, \chi_k \in \hat{G}} \chi_1^{-1}(a_1 b_{\pi(1)}) \cdots \chi_k^{-1}(a_k b_{\pi(k)}) \det((\chi_i(a_j))_{1 \leq i, j \leq k}) \det((\chi_i(b_j))_{1 \leq i, j \leq k}).$$

Observe that

$$\begin{aligned} \Sigma &= \sum_{\chi_1, \dots, \chi_k \in \hat{G}} \sum_{\sigma \in S_k} \text{sgn}(\sigma) \prod_{i=1}^k \chi_i(a_{\sigma(i)} a_i^{-1}) \sum_{\tau \in S_k} \text{sgn}(\tau \sigma) \prod_{i=1}^k \chi_i(b_{\tau \sigma(i)} b_{\pi(i)}^{-1}) \\ &= \sum_{\tau \in S_k} \text{sgn}(\tau) \sum_{\sigma \in S_k} \prod_{i=1}^k \sum_{\chi_i \in \hat{G}} \chi_i(a_{\sigma(i)} b_{\tau \sigma(i)} (a_i b_{\pi(i)})^{-1}) \\ &= \sum_{\tau \in S_k} \text{sgn}(\tau) \sum_{\substack{\sigma \in S_k \\ a_{\sigma(i)} b_{\tau \sigma(i)} = a_i b_{\pi(i)} \text{ for } i=1, \dots, k}} (|G|1)^k. \end{aligned}$$

If $\tau \in S_k$ and $\{a_1 b_{\tau(1)}, \dots, a_k b_{\tau(k)}\} = C$, then τ is identical to π by the assumption of Theorem 1.11(ii), and the identity permutation I is the unique $\sigma \in S_k$ such that $a_{\sigma(i)} b_{\tau \sigma(i)} = a_i b_{\pi(i)}$ for all $i = 1, \dots, k$. Therefore

$$\Sigma = \text{sgn}(\pi) (|G|1)^k \neq 0.$$

So there are $\chi_1, \dots, \chi_k \in \hat{G}$ such that

$$\det((\chi_i(a_j))_{1 \leq i, j \leq k}) \det((\chi_i(b_j))_{1 \leq i, j \leq k}) \neq 0$$

and hence neither $\det((\chi_i(a_j))_{1 \leq i, j \leq k})$ nor $\det((\chi_i(b_j))_{1 \leq i, j \leq k})$ vanishes.

The remaining results in Theorem 1.11(ii) can be easily proved in a similar way.

Acknowledgment. The authors are indebted to the referee for helpful comments.

REFERENCES

- [1] N. Alon, *Combinatorial Nullstellensatz*, *Combin. Probab. Comput.* **8**(1999), 7–29.
- [2] N. Alon, *Additive Latin transversals*, *Israel J. Math.* **117**(2000), 125–130.
- [3] L. Babai, P. Frankl, *Linear Algebra Methods in Combinatorics with Applications to Geometry and Computer Science*, Preliminary Version 2, University of Chicago, 1992.
- [4] S. Dasgupta, G. Károlyi, O. Serra, B. Szegedy, *Transversals of additive Latin squares*, *Israel J. Math.* **126**(2001), 17–28.
- [5] W. D. Gao, D. J. Wang, *Additive Latin transversals and group rings*, *Israel J. Math.* **140**(2004), 375–380.

- [6] W. D. Gao, A. Geroldinger and F. Halter-Koch, *Group algebras of finite abelian groups and their applications to combinatorial problems*, Rocky Mountain J. Math. **39**(2009), 805–823.
- [7] M. Hall, Jr., *A combinatorial problem on abelian groups*, Proc. Amer. Math. Soc. **3**(1952), 584–587.
- [8] D. G. Northcott, *Multilinear Algebra*, Cambridge University Press, Cambridge, 1984.
- [9] H. S. Snevily, *The Cayley addition table of Z_n* , Amer. Math. Monthly **106**(1999), 584–585.
- [10] P. Stevenhagen, H. W. Lenstra, Jr., *Chebotařev and his density theorem*, Math. Intelligencer, **18**(1996), 26–37.
- [11] Z. W. Sun, *Covering the integers by arithmetic sequences. II*, Trans. Amer. Math. Soc. **348**(1996), 4279–4320.
- [12] Z. W. Sun, *On the function $w(x) = |\{1 \leq s \leq k : x \equiv a_s \pmod{n_s}\}|$* , Combinatorica **23**(2003), 681–691.
- [13] Z. W. Sun, *On Snevily’s conjecture and restricted sumsets*, J. Combin. Theory Ser. A **103**(2003), 291–304.
- [14] Z. W. Sun, *An additive theorem and restricted sumsets*, Math. Res. Lett. **15**(2008), 1263–1276.
- [15] T. Tao, *An uncertainty principle for cyclic groups of prime order*, Math. Res. Lett. **12**(2005), 121–127.

DIVISION OF MATHEMATICAL SCIENCES, SCHOOL OF PHYSICAL AND MATHEMATICAL SCIENCES,
NANYANG TECHNOLOGICAL UNIVERSITY, SINGAPORE 637616

E-mail address: FengTao@ntu.edu.sg

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE’S REPUBLIC
OF CHINA

E-mail address: zwsun@nju.edu.cn

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF DELAWARE, NEWARK, DE 19716,
USA

E-mail address: xiang@math.udel.edu